

Instructor: *P. Kumar*

PHYSICS DEPARTMENT

PHY 2049

Exam I

June 1, 2009

Name (print, last first): _____ Signature: _____

*On my honor, I have neither given nor received unauthorized aid on this examination.***YOUR TEST NUMBER IS THE 5-DIGIT NUMBER AT THE TOP OF EACH PAGE.****DIRECTIONS**

- (1) **Code your test number on your green answer sheet (use 76–80 for the 5-digit number).** Code your name on your answer sheet. **Darken circles completely (errors can occur if too light).** Code your student number on your answer sheet.
- (2) **Blacken the circle of your intended answer completely, using a number 2 pencil.** Do not make any stray marks or the answer sheet may not read properly.
- (3) The answers are rounded off. Choose the closest to exact. There is no penalty for guessing.

>>>>>>>**WHEN YOU FINISH**<<<<<<<<<
Hand in the green answer sheet separately.

1. The total negative charge on the electrons in 1 mol of helium (atomic number 2, molar mass 4 g/mole) is:

- (1) $1.9 \times 10^5 \text{C}$ (2) $4.8 \times 10^4 \text{C}$ (3) $9.6 \times 10^4 \text{C}$ (4) $3.8 \times 10^5 \text{C}$ (5) $7.7 \times 10^5 \text{C}$

2. Two identical conducting spheres A and B carry equal charge. They are separated by a distance much larger than their diameters. A third identical conducting sphere C is uncharged. Sphere C is first touched to A, then to B, and finally removed. As a result, the electrostatic force between A and B, which was originally F , becomes:


- (1) $3F/8$ (2) $F/2$ (3) $F/4$ (4) $F/16$ (5) 0

3. Particles 1, with charge q_1 , and 2, with charge q_2 , are on the x axis, with particle 1 at $x = a$ and particle 2 at $x = -2a$. For the net force on a third charged particle, at the origin, to be zero, q_1 and q_2 must be related by $q_2 =$:

- (1) $4q_1$ (2) $2q_1$ (3) $-2q_1$ (4) $-4q_1$ (5) $\frac{-q_1}{4}$

4. A charged oil drop with a mass of 2×10^{-4} kg is held suspended by a downward electric field of 300N/C. The charge on the drop is: ($g = 9.80 \text{ m/s}^2$)


- (1) $-6.5 \times 10^{-6} \text{C}$ (2) $+1.5 \times 10^{-6} \text{C}$ (3) $-1.5 \times 10^{-6} \text{C}$ (4) $+6.5 \times 10^{-6} \text{C}$ (5) 0

5. Two charged particles are arranged as shown. In which region could a third particle, with charge $+1 \text{ C}$, be placed so that the net electrostatic force on it is zero?

- (1) I only (2) I and II only (3) III only (4) I and III only (5) II only

6. Positive charge $+Q$ is uniformly distributed on the upper half a semi-circular rod and negative charge $-Q$ is uniformly distributed on the lower half. What is the direction of the electric field at point P, the center of the semicircle?

- (1) \downarrow (2) \uparrow (3) \leftarrow (4) \rightarrow (5) \nearrow

7. A hollow conductor is positively charged. A small uncharged metal ball is lowered by a silk thread through a small opening in the top of the conductor and allowed to touch its inner surface. After the ball is removed, it will have:
- no appreciable charge
 - a positive charge
 - a negative charge
 - a charge whose sign depends on what part of the inner surface it touched
 - a charge whose sign depends on where the small hole is located in the conductor
8. Positive charge Q is placed on a conducting spherical shell with inner radius R_1 and outer radius R_2 . A particle with charge q is placed at the center of the cavity. The magnitude of the electric field at a point in the shell of the cavity, a distance r from the center, is:
- zero
 - $Q/4\pi\epsilon_0 R_1^2$
 - $q/4\pi\epsilon_0 r^2$
 - $(q + Q)/4\pi\epsilon_0 r^2$
 - $(q + Q)/4\pi\epsilon_0 (R_1^2 - r^2)$
9. A point particle with charge q is at the center of a Gaussian surface in the form of a cube. The electric flux through any one face of the cube is:
- $q/6\epsilon_0$
 - q/ϵ_0
 - $q/4\pi\epsilon_0$
 - $q/3\epsilon_0$
 - $q/12\epsilon_0$
10. What is the net electric potential at point P due to the four particles, if $V = 1$ mV at infinity, $q = 7.00$ fC, and $d = 2.00$ cm?
- 
- 2.6 mV
 - 1.6 mV
 - 3.2 mV
 - 1.6 mV
 - none of these
11. A 140 pF capacitor is charged to a potential difference of 60 V, and the charging battery is disconnected. The capacitor is then connected in parallel with a second (initially uncharged) capacitor. If the potential difference across the first capacitor drops to 48 V, what is the capacitance of this second capacitor?
- 35 pF
 - 20 pF
 - 10 pF
 - 15 pF
 - none of these
12. The electric potential at points in an xy plane is given by $V = (3.0\text{V/m}^2)x^2 - (4.0\text{V/m}^2)y^2$. What is the electric field at the point (3.0 m, 3.0 m)? (E_x, E_y in V/m)
- (-18,24)
 - (18,-24)
 - (27,-36)
 - (-27,36)
 - none of these
13. A nichrome wire is 1m long and $1 \times 10^{-6}\text{m}^2$ in cross-sectional area. When connected to a potential difference of 2V, a current of 4A exists in the wire. The resistivity of this nichrome is:
- $5 \times 10^{-7}\Omega\cdot\text{m}$
 - $2 \times 10^{-7}\Omega\cdot\text{m}$
 - $4 \times 10^{-7}\Omega\cdot\text{m}$
 - $8 \times 10^{-7}\Omega\cdot\text{m}$
 - $10^{-7}\Omega\cdot\text{m}$
14. A flat iron is marked "120V, 600W". In normal use, the current in it is:
- 5A
 - 4A
 - 2A
 - 7.2A
 - 0.2A