

Time-Dependent Perturbation Theory: The Photoelectric Effect

- This handout mirrors the treatment of the photoelectric effect on Shankar pp. 499–506, with two principal differences: (1) The perturbing Hamiltonian is written $H_{1E} = e\mathbf{E} \cdot \mathbf{R}$ instead of $H_{1A} = (e/mc)\mathbf{A} \cdot \mathbf{P}$. (2) The system is assumed to occupy a cubic box of sides L , whereas Shankar treats an infinite system. We comment on the significance of these differences at the end.
- The initial state is taken to belong to the innermost (or K) shell of a hydrogen-like atom of effective nuclear charge Ze , with wave function $\langle \mathbf{r} | i \rangle = \pi^{-1/2} (Z/a_0)^{3/2} \exp(-Z|\mathbf{r}|/a_0)$, where $a_0 = \hbar^2/mc^2$ is the Bohr radius. This state has energy $\varepsilon_i = -Z^2 e^2 / 2a_0 = -(Z\alpha)^2 mc^2 / 2$, $\alpha = e^2/\hbar c$ being the fine-structure constant. The characteristic size of the orbital is $r_0 = a_0/Z = \hbar/(Z\alpha mc)$.

We consider a monochromatic electromagnetic plane wave, $\mathbf{E}(\mathbf{r}, t) = \mathbf{E}_0 \cos(\mathbf{k} \cdot \mathbf{r} - \omega t)$. The electric dipole approximation is valid provided that $|\mathbf{k}|r_0 \ll 1$, or equivalently, $\hbar\omega \ll (Z\alpha)mc^2$. We will consider frequencies in the window $(Z\alpha)^2 mc^2 \ll \hbar\omega \ll (Z\alpha)mc^2$, where not only can we make the dipole approximation, but the final-state energy is sufficiently high that the final state should be well-described by a plane wave of the form $\langle \mathbf{r} | f \rangle = L^{-3/2} \exp(i\mathbf{p}_f \cdot \mathbf{r}/\hbar)$ having an energy $\varepsilon_f = |\mathbf{p}_f|^2/2m$. (See Shankar p. 500 and the end of this handout for discussion of this plane-wave approximation.)

- In the dipole approximation, we need to calculate the dipole matrix element

$$\begin{aligned} \mathbf{r}_{fi} &= \langle f | \mathbf{R} | i \rangle = A \int d^3\mathbf{r} e^{-i\mathbf{p}_f \cdot \mathbf{r}/\hbar} \mathbf{r} e^{-Z|\mathbf{r}|/a_0} \\ &= A \times i\hbar \frac{\partial}{\partial \mathbf{p}_f} \int d^3\mathbf{r} e^{-i\mathbf{p}_f \cdot \mathbf{r}/\hbar} e^{-Z|\mathbf{r}|/a_0} = i\hbar \frac{\partial}{\partial \mathbf{p}_f} \langle f | i \rangle, \end{aligned}$$

where $A = \pi^{-1/2} (Z/La_0)^{3/2}$.

The overlap integral is straightforward to evaluate (see Shankar p. 504):

$$\langle f | i \rangle = A \int d^3\mathbf{r} e^{-i\mathbf{p}_f \cdot \mathbf{r}/\hbar} e^{-Z|\mathbf{r}|/a_0} = \frac{8\pi AZ/a_0}{[(Z/a_0)^2 + (p_f/\hbar)^2]^2}.$$

Therefore

$$\mathbf{r}_{fi} = i\hbar \frac{8\pi AZ/a_0}{[(Z/a_0)^2 + (p_f/\hbar)^2]^3} \left(\frac{-4\mathbf{p}_f}{\hbar^2} \right).$$

Noting that $(Z/a_0)^2 + (p_f/\hbar)^2 = 2m(\varepsilon_f - \varepsilon_i)/\hbar^2$, we find

$$\mathbf{r}_{fi} = -\frac{2i\hbar}{m(\varepsilon_f - \varepsilon_i)} \mathbf{p}_f \langle f | i \rangle = -i \frac{4\pi AZ\hbar^5}{a_0 m^3 (\varepsilon_f - \varepsilon_i)^3} \mathbf{p}_f. \quad (1)$$

- Fermi's Golden Rule gives the scattering rate from $|i\rangle$ to $|f\rangle$ as

$$R_{i \rightarrow f} = \frac{2\pi}{\hbar} \left| \frac{e}{2} \mathbf{E}_0 \cdot \mathbf{r}_{fi} \right|^2 \delta(\varepsilon_f - \varepsilon_i - \hbar\omega).$$

In order to calculate the differential scattering cross-section $d\sigma/d\Omega$, defined by

$$\frac{d\sigma}{d\Omega} = \frac{\text{power absorbed by atom while emitting electrons into solid angle } d\Omega}{(\text{incident energy flux of electromagnetic field}) \times d\Omega},$$

we need to find the density of final states \mathbf{p}_f . If we apply periodic boundary conditions to the cubic box, then the allowed final states obey

$$(p_f)_j = \hbar \frac{2\pi n_j}{L} = \frac{\hbar n_j}{L} \quad \text{for } j = x, y, z.$$

Thus, the number of allowed states in a momentum-space volume element $p_f^2 dp_f d\Omega$ is

$$\left(\frac{L}{\hbar}\right)^3 p_f^2 dp_f d\Omega = \left(\frac{L}{\hbar}\right)^3 m p_f d\varepsilon_f d\Omega,$$

and the power absorbed by the atom in scattering into solid angle $d\Omega$ is

$$P_{i \rightarrow d\Omega} = \hbar\omega R_{i \rightarrow d\Omega} = d\Omega \int d\varepsilon_f \left(\frac{L}{\hbar}\right)^3 m p_f R_{i \rightarrow f} = \frac{e^2}{\pi} \left(\frac{Z}{a_0}\right)^5 \frac{p_f^3 E_0^2}{m^5 \omega^6} |\hat{\mathbf{E}}_0 \cdot \hat{\mathbf{p}}_f|^2 d\Omega.$$

The incident energy flux of the electromagnetic wave is $J_{\text{in}} = uc = (c/4\pi) |\mathbf{E}(\mathbf{r}, t)|^2$, or, averaged over one complete cycle, $J_{\text{in}} = (c/8\pi) |\mathbf{E}_0|^2$. Thus,

$$\frac{d\sigma}{d\Omega} = \frac{P_{i \rightarrow d\Omega}}{J_{\text{in}} d\Omega} = \frac{8e^2 \hbar p_f^3}{c} \left(\frac{Z}{m\omega a_0}\right)^5 |\hat{\mathbf{E}}_0 \cdot \hat{\mathbf{p}}_f|^2. \quad (2)$$

- **Comment 1:** The finite system size does not enter the final result. If one works with an infinite system, the correct density of final states is ensured through the delta-function normalization of the plane wave, i.e., $\langle \mathbf{r} | \mathbf{p} \rangle = (2\pi\hbar)^{-3/2} \exp(i\mathbf{p} \cdot \mathbf{r}/\hbar)$.
- **Comment 2:** Equation (2) agrees, for instance, with that obtained from H_{1E} in *Quantum Theory of Light* by R. Loudon (Clarendon Press, Oxford, 1973). On the other hand, this $d\sigma/d\Omega$ is 4 times greater than that given by Shankar, Merzbacher, and Sakurai, all of which authors use H_{1A} .

This discrepancy appears to stem from the fact that the general result (see Shankar p. 503)

$$\langle f | \mathbf{P} | i \rangle = \frac{im(\varepsilon_f - \varepsilon_i)}{\hbar} \langle f | \mathbf{R} | i \rangle$$

holds only if $|i\rangle$ and $|f\rangle$ are *exact* eigenstates of the same Hamiltonian. Here, $|f\rangle$ is only an approximate version of the true final state, which is a plane wave plus an incoming spherical wave (see Merzbacher p. 502). Since $|f\rangle$ is not a true eigenstate of the Coulomb Hamiltonian, different formulations of the dipole approximation are not guaranteed to produce the same result. It appears that \mathbf{d}_{fi} given in Eq. (1) is twice the correct value. It can be shown (e.g., see Ch. 12 of *Intermediate Quantum Mechanics* by H. A. Bethe and R. Jackiw (2nd Edition, W. A. Benjamin, Reading, Massachusetts, 1968) that using H_{1E} with the exact final state wave functions yields the result obtained using H_{1A} with the plane-wave approximate wave functions. *This should be a salutary warning that calculating the effects of radiation on matter is a subtle business!*