

## A Short Simple Evaluation of Expressions of the DebyeWaller Form

N. David Mermin

Citation: *J. Math. Phys.* **7**, 1038 (1966); doi: 10.1063/1.1704995

View online: <http://dx.doi.org/10.1063/1.1704995>

View Table of Contents: <http://jmp.aip.org/resource/1/JMAPAQ/v7/i6>

Published by the [American Institute of Physics](#).

---

### Additional information on J. Math. Phys.

Journal Homepage: <http://jmp.aip.org/>

Journal Information: [http://jmp.aip.org/about/about\\_the\\_journal](http://jmp.aip.org/about/about_the_journal)

Top downloads: [http://jmp.aip.org/features/most\\_downloaded](http://jmp.aip.org/features/most_downloaded)

Information for Authors: <http://jmp.aip.org/authors>


## ADVERTISEMENT

The most comprehensive support for physics in any mathematical software package  
World-leading tools for performing calculations in theoretical physics

**Maple™ 16**  
The Essential Tool for Mathematics and Modeling  
[www.maplesoft.com/physics](http://www.maplesoft.com/physics)

- Your work in Maple matches how you would write the problems and solutions by hand
- State-of-the-art environment for algebraic computations in physics
- The only system with the ability to handle a wide range of physics computations as well as pencil-and-paper style input and textbook-quality display of results
- Access to Maple's full mathematical power, programming language, visualization routines, and document creation tools

*InertiaTensor* :=  $\sum_{k=1}^n m_k \left( \|\vec{r}_k\|^2 \right)$   
 $\Gamma_{2,2}^1 = \frac{(1 + 2)}{\partial r}$



## A Short Simple Evaluation of Expressions of the Debye-Waller Form

N. DAVID MERMIN

*Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York*

(Received 15 November 1965)

Averages like those encountered in the theory of the Debye-Waller factor are evaluated in one sentence.

WHEN calculating absorption, emission, or scattering cross sections for crystalline matter in the harmonic approximation one needs the thermal equilibrium average of exponentials of operators linear in the atomic displacements and/or momenta:

$$\langle e^{\sum c_i a_i + d_i a_i^\dagger} \rangle = \text{Tr} e^{-\beta H} e^{\sum c_i a_i + d_i a_i^\dagger} / \text{Tr} e^{-\beta H},$$

$$H = \sum \omega_i (a_i^\dagger a_i + \frac{1}{2}), \quad \beta = 1/k_B T, \quad [a_i, a_j^\dagger] = \delta_{ij}. \tag{1}$$

This can be evaluated in a variety of ways,<sup>1</sup> some difficult, some direct, but all annoyingly cumbersome considering the simplicity of the final form. Here is a derivation as simple as the result:

As in most approaches, begin by using the well-known formula<sup>1</sup>

$$e^{A+B} = e^A e^B e^{-\frac{1}{2}[A,B]} \quad ([A, B] \text{ a } c\text{-number}) \tag{2}$$

to reduce (1) to

$$\langle e^{\sum c_i a_i + d_i a_i^\dagger} \rangle = \langle e^{\sum c_i a_i} e^{\sum d_i a_i^\dagger} \rangle e^{-\frac{1}{2} \sum c_i d_i}, \tag{3}$$

but instead of proceeding with the clumsy direct evaluation of

$$g(c_i, d_i) = \langle e^{\sum c_i a_i} e^{\sum d_i a_i^\dagger} \rangle, \tag{4}$$

note that (2) also entitles one to conclude

$$\langle e^{\sum c_i a_i + d_i a_i^\dagger} \rangle = \langle e^{\sum d_i a_i^\dagger} e^{\sum c_i a_i} \rangle e^{\frac{1}{2} \sum c_i d_i}, \tag{5}$$

which is consistent with (3) only if

<sup>1</sup>See, for instance, A. A. Maradudin, E. W. Montroll, and G. H. Weiss, *Solid State Phys. Suppl.* **3**, 239 (1963).

$$\begin{aligned} g(c_i, d_i) &= e^{\sum c_i d_i} \langle e^{\sum d_i a_i^\dagger} e^{\sum c_i a_i} \rangle \\ &= e^{\sum c_i d_i} \langle e^{\beta H} e^{\sum c_i a_i} e^{-\beta H} e^{\sum d_i a_i^\dagger} \rangle \\ &\quad \text{(cyclical permutation within trace)} \\ &= e^{\sum c_i d_i} \langle e^{\sum c_i e^{\beta H} a_i e^{-\beta H}} e^{\sum d_i a_i^\dagger} \rangle \\ &= e^{\sum c_i d_i} \langle e^{\sum c_i e^{-\beta \omega_i} a_i} e^{\sum d_i a_i^\dagger} \rangle \\ &\quad \text{(harmonic approximation)} \\ &= e^{\sum c_i d_i} g(c_i e^{-\beta \omega_i}, d_i), \end{aligned} \tag{6}$$

from which identity it follows at once (by iteration or induction on  $n$ ) that

$$g(c_i, d_i) = e^{\sum c_i d_i (1 + e^{-\beta \omega_i} + \dots + e^{-n\beta \omega_i})} g(c_i e^{-(n+1)\beta \omega_i}, d_i), \tag{7}$$

and hence, taking the limit  $n \rightarrow \infty$  (each  $\omega_i$  is positive),

$$g(c_i, d_i) = e^{\sum c_i d_i (1 - e^{-\beta \omega_i})^{-1}} g(0, d_i), \tag{8}$$

which, since it follows trivially from (4) that

$$g(0, d_i) = \langle e^{\sum d_i a_i^\dagger} \rangle = 1, \tag{9}$$

completes the derivation:

$$\begin{aligned} \langle e^{\sum c_i a_i + d_i a_i^\dagger} \rangle &= e^{-\frac{1}{2} \sum c_i d_i} g(c_i, d_i) \\ &= e^{\frac{1}{2} \sum c_i d_i \coth \frac{1}{2} \beta \omega_i}. \end{aligned} \tag{10}$$